

Strength distribution of solar magnetic fields in photospheric quiet Sun regions

J.C. RAMÍREZ VÉLEZ¹, A. LÓPEZ ARISTE² and M. SEMEL¹

¹ LESIA, Observatoire de Paris Meudon, 92195 Meudon, France; e-mail: Julio.Ramirez@obspm.fr, Meir.Semel@obspm.fr

² THEMIS, CNRS UPS 853, c/vía Láctea s/n. 38200. La Laguna, Tenerife, Spain; e-mail: arturo@themis.iac.es

Accepted in A&A

ABSTRACT

Context. The magnetic topology of the solar photosphere in its quietest regions is hidden by the difficulties to disentangle magnetic flux through the resolution element from the field strength of unresolved structures. The observation of spectral lines with strong coupling with hyperfine structure, like the observed Mn I line at 553.7 nm, allows such differentiation.

Aims. To analyse the distribution of field strengths in the network and intranetwork of the solar photosphere through inversion of the Mn I line at 553.7 nm.

Methods. An inversion code for the magnetic field using the Principal Component Analysis (PCA) has been developed. Statistical tests are run on the code to validate it. The code has to draw information from the small-amplitude spectral feature appearing in the core of the Stokes V profile of the observed line for field strengths below a certain threshold, coinciding with lower limit of the Paschen-Back effect in the fine structure of the involved atomic levels.

Results. The inversion of the observed profiles, using the circular polarization (V) and the intensity (I), shows the presence of magnetic fields strengths in a range from 0 to 2 kG, with predominant weak strength values. Mixed regions with mean strength field values of 1130 and 435 Gauss are found associated with the network and intranetwork respectively.

Conclusions. The Mn I line at 553 nm probes the field strength distribution in the quiet sun and shows the predominance of weak, hectoGauss fields in the intranetwork, and strong, kiloGauss fields in the network. It also shows that both network and intranetwork are to be understood at our present spatial resolutions as field distributions of which we hint the mean properties.

Key words. Sun: magnetic fields - Sun: photosphere - Line: profiles - Methods: data analysis

1. Introduction

The absence of observables of the magnetism of the quiet sun is the main handicap in the determination of the magnetic topology of this region covering most of the solar photosphere. Over-interpretation of the scarce information contained in the usually observed spectral lines in magnetometry, and its comparison with numerical simulations of magnetoconvection, has been the subject of long disputes about the nature of those fields. The addition of further observables, able to constrain the models and to unveil hidden ambiguities and biases in the diagnostic techniques was mandatory in such a situation. This work is a contribution in that direction, with a further step forward in the use of Mn lines with a strong hyperfine coupling.

In a short descriptive summary of the previous studies of the magnetism in quiet Sun regions, we find from one side, the Fe lines in the visible region whose circular polarization amplitudes are almost uniquely sensitive to magnetic flux, (e.g. Keller et al. 1994; Sánchez Almeida & Lites 2000; Lites 2002, Domínguez Cerdeña et al. 2003; Orozco Suárez et al. 2007), and on the other side, we find the inversions of the Fe lines in the near-IR domain whose profiles are almost universally split by Zeeman effect (Lin 1995; Lin & Rimmele 1999; Khomenko et al. 2003). Traditionally, studies based on the visible spectral lines concluded on the presence of fields with kG strengths and small surface coverage, while studies based on infrared lines favoured weaker hG fields and more spread in surface.

Recently, the picture has been made more realistic by considering a continuum distribution function for magnetic field strengths at each resolution element in the quiet sun (Socas Navarro & Sánchez Almeida 2003), instead of a single magnetic vector value. Additionally, a very interesting new approach has enriched the individual line analysis in quiet regions: the inversions of simultaneous and co-spatial observations of the Fe lines in the visible (630 nm) and in the near-IR (1.5 μ m). Two such studies have already been carried out founding divergent results. In the first of these works, Domínguez Cerdeña et al. (2006) have employed a model with three magnetic components (MISMAS model, Sánchez Almeida & Landi Degl'Innocenti 1996) to simultaneously invert the lines in both spectral ranges. The authors retrieved field distributions in the second and third magnetic components with peaks in the strong regime and they conclude that the kG fields contribution dominates the magnetic flux and energy transport. In the other of these studies, Martínez González et al. (2008) showed that when only the inversions of the Fe lines in the visible are considered, the magnetic distribution is dominated by strong strength fields (kG). However, when simultaneously near-IR and visible inversions were both considered, the kG contributions disappears and the distributions are mostly dominated by weak strength fields (hG). This last conclusion is in accord to the results obtained from near-IR inversions alone (see previous references). These results, suggesting a probable bias of the quiet sun Fe inversions when performed only in the visible at 630 nm had also been found on numerical tests (Bellot Rubio & Collados 2003;

Martínez González et al. 2006). In the counterpart, a possible bias in the inversions of the IR alone, has been also argued suggesting that this lines are slightly sensitive to the strong strength fields, such that when mixed and not resolved magnetic structures of hG and kG are both present, the inferred strengths of hG are preferred over those of kG (e.g. Sánchez Almeida & Lites 2000; Domínguez Cerdeña et al. 2006). Such discussions and differing results illustrate the difficulty and ambiguity of the measurements attempted, and have prompted the search for more observables.

Apart from those Zeeman-based techniques discussed above, the description should be completed with the Hanle effect diagnostics, pioneered by Stenflo (1982). A paradigm of not structured and turbulent fields, e.g. (Manso Sainz et al. 2004), could result in apparent contradiction with the tube-like structures used to describe Zeeman-based observations if one insists in the picture of a single vector magnetic field per resolution element.

Mn lines subject to strong hyperfine coupling appeared as an interesting observable to be added to the previous ones: their amplitudes are dependent on magnetic flux as the usual Fe lines in the visible regime, but under certain magnetic field regimes a new spectral feature appears in the Mn profile. Just the presence or absence of such a spectral feature allows the observer to determine the presence or absence of fields with strengths below a certain threshold thus adding a new observable on field strength to be compared to measurements with near-IR lines.

Quiet sun conditions limit the information on magnetic fields to circular polarization profiles, whose amplitudes stay too near to the usual signal-to-noise levels found in solar polarimetry. The amplitudes of the linear polarization Stokes parameters use to be below the noise level at the typical spatial resolutions, and are thus useless for analysis purposes; and intensity profiles are insensitive to the weak magnetic fluxes involved. Consequently one does not only need to add further spectral lines to the amount of observables, but also do it under observing conditions that guarantee very high signal-to-noise levels in polarimetry. We used the telescope THEMIS for that purpose in this work and we explored the capabilities of the Mn lines to provide further information on the quiet sun magnetic strength distribution. We simplified the formation of the Mn lines with a radiative transfer model using a Milne-Eddington atmosphere; the model proves to be at the limit of present possibilities with yet too many of its parameters left undetermined by the available observables. We nevertheless prove through synthetic tests that we can ascertain the distribution of field strengths in strong and weak flux regions (photospheric network and intranetwork respectively) with good accuracy, within the limits of the adopted Milne-Eddington modeling for the Sun.

With the results presented here, that show a field strength distribution with maximum number of occurrences at very weak fields (less than or around 100 G), we propose in the conclusions a cartoon scenario that would be consistent with both the Zeeman-based measurements (visible and near-IR) and the Hanle measurements.

In section 2, we review the signatures of the magnetic field strength in the line formation of the Mn I profiles. In section 3, we describe the inversion code used in this work to retrieve the strength and magnetic flux from the data, and we test its capabilities with realistic noise conditions. The observations are described in section 4 and the results presented in section 5. Conclusions and a final overview are included in the final section.

2. Hyperfine structure and the Stokes parameters

The coupling of the nuclear angular momentum (I) with the orbital total angular momentum (J_{LS}) gives rise to hyperfine structure (HFS). The total angular momentum $F = J + I$ results in new non-degenerated atomic levels. The apparent splitting of the atomic transitions due to this effect, and the consequent broadening of solar and stellar spectral lines has been discussed in several papers since the work of Abt (1952), and used to measure the abundances of different elements (Kurucz 1993).

The most common observable for the analysis of (solar or stellar) spectral lines modified by hyperfine structure is the broadening of the line profile. Such broadening however can be the result of others physical phenomena as well: micro-turbulence, Doppler broadening or just blended lines. Too often the ambiguity on the origin of an observed line broadening cannot be solved for, even in observations with very high spectral resolution, unless full spectropolarimetry is performed. Such is the case presented in this work, the Mn I line at 553.7 nm: although producing a slightly asymmetric intensity profile, the recognition of the HFS features becomes straightforward in circular polarisation, since the HFS together with an atmosphere permeated of an external magnetic field produce a unique spectral signature in the Stokes V profiles. The use of this signature and its variations under different conditions as a tool for the detection of magnetic fields in the quiet sun in different field strength regimes has been theoretically first proposed in López Ariste et. al (2002). Later, in López Ariste et. al (2006), the authors showed that observationally it is possible to detect the HFS effect in the circular polarization profiles. In this work we extend previous studies and we build an inversion code drawing its information from those spectral features in the Stokes profiles due to HFS. The inversion algorithm is firstly validated and characterized with synthetic data, and then applied to observed data. A quantitative analysis of the field strength distribution in the quiet sun results.

For illustration and reference we show, in Fig. 1, the calculated spectral signature of the Stokes V profiles for the Mn I at 553.7 nm, as a function of the magnetic field strength. The important remark from the V profiles pictured is that the weaker the magnetic field strength, the higher the amplitude of the spectral feature due to the HFS, visible in the central part of the profile.

3. The inversion code

3.1. Solution to the radiative transfer equations

The four coupled equations for the radiative transfer of the Stokes parameters were solved analytically by the code `DIAGONAL` (López Ariste & Semel 1999), which was slightly modified from its original version to include the computation of hyperfine structure.

To use the code `DIAGONAL` we assume a Milne-Eddington atmosphere, with the possibility to include numerically gradients with optical depth of the velocity and magnetic fields, and adding up to fifteen free parameters. Evidently, the use of a bigger number of free parameters implies a higher risk of inversion ambiguities and biases, if they happen to exceed the dimensionality of the observables. But on the other hand considering less parameters may not allow to cover all the variety of the observed profiles. We decided thus to limit the number of parameters for each atmospheric model, to the ones described below, leaving out the velocity and magnetic gradients along the line of sight.

The inversion code works in hierarchical mode by a double pass fit. In the first pass only the intensity is fitted, permitting




Fig. 1. Stokes V profiles as a function of the magnetic field strength. Note that the spectral feature on the central part of the profiles is the signature associated to the HFS regime, and that it disappears about $B \sim 750$ Gauss, as the line passes into the incomplete Paschen-Back regime.

half of the atmospheric parameters (Doppler width, V_D , velocity in the line of sight V_{LOS} , η_0 and the source function gradient, $\nabla\tau$) to be constrained in their ranges of variation. The remaining four atmospheric parameters are related to the presence of a magnetic field: the angle between the magnetic field direction and the direction of the line of sight θ_{BLOS} , the azimuthal angle θ_z , the strength of the magnetic field B and a filling factor ff , accounting for unresolved structure. In the second (and final) fit, the eight parameters are considered in fitting simultaneously the profiles of intensity (I) and circular polarization (V). Each atmospheric parameter will be left to vary randomly in the range of values specified in table 1 in order to build a database for the inversions.

3.2. Atmospheric models and database

Let χ denote a given combination of the described atmospheric parameters such that,

$$\chi_m = (V_D, V_{LOS}, \eta_0, \nabla\tau, B, \theta_{BLOS}, \theta_z, ff)_m \quad (1)$$

is the m_{th} combination. Let $S_m(\chi_m)$ denote the Stokes vector calculated for this combination of atmospheric parameters χ_m :

$$S_m(\chi_m) = (I_m(\chi_m), Q_m(\chi_m), U_m(\chi_m), V_m(\chi_m)), \quad (2)$$

and let P_m denote any of the components of $S_m(\chi_m)$.

In the computation Stokes parameters we have considered 201 spectral wavelength points with a step size of $12 \text{ m}\text{\AA}$, covering in total $2.41 \text{ \AA}$ around the line center at 553.762 nm . In the following, we proceed to consider each profile P_m as a vector of 201 points P_m . The creation of the database is the most important step for the inversions working with principal components analysis (e.g. Rees et al. 2000; Socas-Navarro et al. 2001). It is required to consider a large enough¹ number of profiles in order to consider the database as statistically representative. Thus, firstly, we have created a database of 80802^1 profiles P_m , hereafter called the profile space matrix ($\mathcal{M}$), and we assume that all

possible observed profiles in the data set are represented in the database. This assumption is however verified in the tests presented at the end of this section.

We next perform a *Singular Value Decomposition* (SVD) (Golub & Van Loan 1996) of the space matrix to compute an algebraic basis of the space spanned by the profiles. We should refer to the elements of the basis as eigenprofiles.

3.3. Analysis with PCA

Following the synthetic notation of Eq. (2), for each one of the Stokes parameters (I, Q, U, V) it is possible to construct the 201×80802 profile matrix $\mathcal{M}$ that contains all the profiles P_m of the database:

$$\mathcal{M} = (P_0 \cdots P_{80801})^T. \quad (3)$$

After computation of the eigenprofiles, each profile in the matrix $\mathcal{M}$ can be reconstructed as:

$$P_m(\chi_m) = \sum_i \alpha_i(\chi_m) \hat{e}_i \quad ; i = 0, 1, \dots, 200 \quad (4)$$

where $\hat{e}_i$ are the eigenvectors associated to the space profile ($\mathcal{M}$) and α_i are the coefficients of the linear combination associated to the profile P_m . Since the eigenvectors form an orthonormal basis of the space ($\mathcal{M}$) arranged in order of importance, one can cut the expansion in Eq. (4) and preserve only the first eigenvectors that contain relevant information on the shape of the profile. The rest of eigenvectors are associated with the noise and there is no interest in preserving them (Hansen 1992; Rees et al. 2000). In our study we find that the appropriate number of components in Eq. (4), is $i_{max} = 30$.

With the help of the set of eigenvectors, $\{\hat{e}_i\}$, we create a second database ($\mathcal{M}'$) that is independent from the ($\mathcal{M}$) matrix, used to compute the $\{\hat{e}_i\}$. The ($\mathcal{M}'$) database will be decomposed in terms of the coefficients α_i as follows. Let us denote P_n a vector of the new (second) database. Using the Eq. (4),

$$\alpha_i(\chi_n) = P_n(\chi_n) \cdot \hat{e}_i \quad ; i = 0, 1, \dots, 29. \quad (5)$$

In this way, each profile P_n (observed or calculated) has associated a unique set of coefficients ($\alpha_0, \dots, \alpha_{i_{max}}$), denoted hereafter $\alpha_n \equiv (\alpha_0, \dots, \alpha_{i_{max}})$. So instead of working in the profile matrix space ($\mathcal{M}'$) it is equivalent, and faster, to work in the coefficient's space ($\mathcal{C}$). From Eq. (3), and in analogy with ($\mathcal{M}'$), we obtain for each profile P_n the correspondent coefficients α_n in order to build the 30×80802 coefficient matrix ($\mathcal{C}$),

$$\mathcal{C} = (\alpha_0 \cdots \alpha_{80801})^T. \quad (6)$$

3.4. Description and validation of the inversion code

One important quality of the code developed is that it is very easy to estimate the error bars per model parameter. We present here the tests applied for that purpose before pursuing into the inversion of the observed profiles.

Let P_{syn} denote a synthetic profile with associated coefficients α_{syn} . The profile P_{inv} , considered to be the solution by the inversion code, is found in the space of coefficients ($\mathcal{C}$) as the nearest set of coefficients (in an Euclidean sense) to α_{syn} :

$$\| \alpha_{inv} - \alpha_{syn} \| = \min(\| \alpha_{syn} - \alpha_n \|) \quad ; n = 0, 1, \dots, 80801. \quad (7)$$

Since each one of the Stokes parameters would give, if considered separately, one solution to the Eq. (7), let us say $(\alpha_{inv,I}, \alpha_{inv,Q}, \alpha_{inv,U}, \alpha_{inv,V})$, it is required to find the best solution

¹ The actual figure has no particular meaning other than illustrating what we consider as *large enough* and for future reference.

Table 1. All the parameters vary randomly in the indicated ranges. The first four parameters of the *initial set*, are used to fit the intensity in the first inversion passage in the code. In the *final set*, all the eighth parameters are considered to invert simultaneously (I) and (V).

Parameters	V_D (mÅ)	η_0	$\nabla\tau$	V_{LOS} (km/s)	B (G)	θ_{BLOS} (deg.)	θ_z (deg.)	ff
<i>Initial Set</i>	[35,75]	[0,2.5]	[0,15]	[-4,4]	[0,2000]	[0,90]	[0,90]	[0,1]
Inverted (max,min)	(41,60)	(0.7,1.8)	(1.0,14.9)	(0,0.54)	—	—	—	—
mean value	47	1.16	8.8	0.02	—	—	—	—
<i>Final Set</i>	[39,61]	[0.5,2]	[0,15]	[-2,2]	[0,2000]	[0,90]	[0,90]	[0,1]

for the four Stokes parameters simultaneously. For this purpose and given that it is a well-known fact, but particularly true in our data sets, that $I_n \gg V_n > U_n, Q_n$ necessarily implying that, in average, $\alpha_{inv,I} > \alpha_{inv,V} > \alpha_{inv,U}, \alpha_{inv,Q}$ (e.g. López Ariste & Casini 2002), we find useful to introduce a *weight* for each solution. Thanks to this, the polarized signals are weighted similarly to the intensity one, despite their disparate amplitudes.

Let the scalars

$$\alpha_{w,S} = \frac{\max(\alpha_{syn,I})}{\max(\alpha_{syn,S})} \quad ; S = I, Q, U, V \quad (8)$$

be the factors that modify the computation of the norms in Eq. (7), such that each one of the four solutions has the appropriate *weight* and can be added to the others in a linear way. The final solution, α_{sol} , is then found as:

$$(\sum_S \alpha_{w,S} \|\alpha_{inv,S} - \alpha_{syn,S}\|) = \min(\sum_S \alpha_{w,S} \|\alpha_{syn,S} - \alpha_{n,S}\|) \quad (9)$$

where S denotes, as usual, the Stokes parameters (I,Q,U,V).

Since we know from observations, described in section 4, that the signal-to-noise ratio in linear polarization (Stokes Q, U) is not big enough to include them in the analysis, they were excluded as well from the tests applied to the code and only the intensity and circular polarization were considered. This implies that from Eq. (8), $S = I, V$.

Finally, we compare the solution values of χ_{sol} (associated to α_{sol}) to the known values of χ_{syn} (associated to α_{syn}). This comparison leads to the statistical determination of the inversion errors per model parameter, where by error we mean the absolute difference between the original and the retrieved values.

3.4.1. Discussion of the tests

In order to not bias the tests we have constrained the thermodynamic parameters to the ranges of variation retrieved from the observational data sets after the first inversion (see the first three rows of table 1). Otherwise, we have not imposed any *a priori* constrain to the magnetic parameters, and the profiles used in the tests were synthesized using independent uniform distributions for each parameter, following the values found in the last row of table 1. However, in the search of direct translation to the inversion of quiet sun profiles, we have conditioned the profiles to have amplitudes in the V profiles that do not exceed the maximum amplitude found in the data set ($V < 5 \times 10^{-3}$) while being bigger than the lower noise level ($V > 5 \times 10^{-5}$). Under these conditions, it is expected that the sample of profiles used in the tests is comparable with that from quiet sun data.

In Fig. 2, we show the results of the inversion of samples of 500 profiles. As a first test, we have not included noise in the inverted profiles. With the exceptions of $\nabla\tau$ and θ_z , the errors in most of the parameters decrease rapidly to zero.

The bad results for θ_z are due to the fact that the linear polarization, Stokes Q, U , was not included in the inversions, implying that the inverted values in this parameter correspond to an




Fig. 2. Inversions histograms of each atmospheric parameter. The number in the upper right corner indicates the value of the bin size used.

aleatory distribution. Consequently, the original (by construction) and the inverted distributions in the azimuthal parameter are aleatory. Finally, plotting the absolute difference between these two distributions results in an histogram with a pick error centered in (around) zero and that decrease smoothly, like the one obtained for θ_z .

Now, the most relevant parameter in our analysis is the magnetic field strength B , since the major interest of this work is to analyse the magnetic field in quiet sun conditions. In what follows we focus the discussion on this parameter and we include also the results of inversion of profiles with noise.

3.4.2. Analysis of effective inversions in B

The graphics in Fig. 2 correspond to the ideal case of profiles without noise. Considering that the real profiles have a typical noise level of the order of 2×10^{-4} to 5×10^{-5} the continuum level, we proceed to invert synthetic profiles with noise added to those levels. Let B_{inv} denote the retrieved value of the field strength, and let B_{syn} denote the real atmospheric value of the synthetic profile. Representing by (ΔB) the upper limit of the error value, such that

$$\Delta B = |B_{inv} - B_{syn}|, \quad (10)$$

we can characterize the inversion efficiency behaviour of the code, understood as the percentage of profiles correctly inverted as a function of (ΔB) . In table 2, we compare the inversions of the magnetic strength for profiles with different noise levels, and in Fig. 3 we plot the respective cumulative distribution functions.

Table 2. Inversions efficiency of the magnetic field strength, expressed in percentage of correct inversions as function of ΔB .

ΔB	Noise level		
	0	5×10^{-5}	2×10^{-4}
50 G	69 %	46 %	40 %
100 G	89 %	74 %	65 %
150 G	96 %	88 %	80 %
200 G	99 %	93 %	89 %
300 G	100 %	98 %	96 %




Fig. 3. Curves of cumulative percentage of correct inversions. Each sample consisted of 500 profiles. The solid line corresponds to profiles without noise. The dashed line corresponds to a noise level of 5×10^{-5} and the dash-point-dash line to a noise level of 2×10^{-4} the intensity of the continuum. The inset indicates the efficiency percentage for an upper limit of $\Delta B = 150$ G.

The value of the magnetic field strength goes from 0 through 2000 G covering all the range of magnetic strengths inherent to the quiet Sun regime. Since the spectral signature in Stokes V profiles evidence of HFS regime, like those in Fig. 1, disappears for values around $B \sim 750$ G, we have special interest in the behaviour of the code while inverting profiles in the

weak field regime. In table 3 we show that the precision inverting the magnetic strength is mainly independent of the magnitude of the magnetic field. The inversion gently improves with higher values of the field strength, but in general the precision in the result is preserved. We conclude that, assuming a Milne-

Table 3. Inversion's efficiency percentages in the weak field regime as function of the magnetic field strength.

ΔB	$B \leq 150G$	$B \leq 350G$	$B \leq 550G$	$B \leq 750G$	Noise
100G	75 %	77 %	81 %	84 %	No
200G	98 %	99 %	99 %	99 %	
100G	60 %	57 %	61 %	64 %	5×10^{-5}
200G	85 %	87 %	89 %	90 %	
100G	56 %	53 %	54 %	57 %	2×10^{-4}
200G	76 %	80 %	82 %	83 %	

Eddington model, the code developed in this work for the line of Mn I at 553.7 nm, correctly inverts profiles in very different field strength regimes: from very weak magnetic fields (lower than 100 G) up to 2 kG, with an error bar of 200G with 89 % of probability (300G with 96 % of probability) independent of the field strength. With such results we can assert that, even if the actual field strength cannot be measured from inversions with sufficient precision, at least the inversion algorithm can disentangle without any ambiguity the hG from the kG regimes of field strength.

4. Observations

We carried out two campaigns of observations at the THEMIS telescope (López Ariste et al. 2000; Gelly et al. 2006) in the multiline observation mode MTR. During the first campaign in June 2005, we searched for the spectral signature of the hyperfine structure of the Mn I atom among four candidate lines in the visible spectral region at wavelengths 539.4 nm, 547.0 nm, 551.6 nm and 553.7 nm, respectively. From the data, we concluded that out of these four lines, only the one at 553.7 nm presented a detectable spectral signature of HFS in the Stokes V profiles clear enough to be used for diagnostics. During the second campaign in June 2006 we focused on this line. In both campaigns the Fe I doublet at 630.1 and 630.2 nm was observed simultaneously with the Mn I lines. Some relevant results concerning the importance of simultaneously observing spectral lines from both atomic species, as well as a brief description of the data acquisition and reduction, has been presented in (López Ariste et al. 2006).

In both campaigns we scanned the solar disk center and, in at least one day per campaign, we enjoyed excellent seeing conditions in the early morning. We have used the polarized signals levels of the Fe doublet (630.2 nm), observed simultaneously and co-spatially, to verify that around the scanned area there were no traces of active regions, nor enhanced magnetic areas: the Stokes V amplitudes of this magnetically sensitive line stay below 1% polarization at all times, indicative of mostly quiet photosphere, both network and intranetwork regions alike. The data sets consisted of temporal sequences of 100 modulation cycles in polarization. Each modulation cycle was made of 6 images with exposure times of either 300 ms or 500 ms. The slit was kept fixed over the solar disk, up to residual seeing-induced image movements. In 2006 the new THEMIS correlation tracker was fully functional, thus further diminishing any residual movement. The total time of these sequences, around 8 minutes, is comparable to the lifetime of the granulation. The dimensions of

the slit spanned a region of 48 arc sec in two separate regions of 15.5 arc secs each, with a masked region of 17 arc sec between them (this masked region is a requirement of the polarimetry mask used for the observation, which places the splitted, orthogonal polarisation, beams 16 arcsec away one from the other in the direction along the slit). The slit width was kept at 0.5 arc sec resulting in a spectral resolution of 20 mÅ in agreement with the pixel spectral sampling of 10 mÅ.

We reduced the data with the *DeepStokes* software tool currently integrated to the THEMIS system facilities, modified to preserve spatial resolution. As delivered the *DeepStokes* code sums all images in the temporal series to increase the signal-to-noise ratio. Because of the long-time integration, image motion and blurring results in a degradation of the image quality. Trying to avoid such a degradation of the spatial resolution, otherwise attained in the individual images of the time series, we modified the code so that the addition of further images in the time series was not done regardless of a further degradation in the image quality.

Specifically, in order to decide if the modulation cycle $i + 1$ was to be added to the sum of the previous i modulation cycles, a cut along the slit (at a pre-specified continuum wavelength of the intensity image) for the modulation cycle $i + 1$ is compared to an analogous cut in the previous added i cycles. The number N of points which differed in both such cuts in less than D was measured. When N happened to be bigger than a pre-specified threshold N_0 the $i + 1$ cycle was deemed *spatially coherent* with the previous ones and its Stokes parameters added to the series, increasing the signal-to-noise ratio. If the $i + 1$ cycle failed to pass the test, it was assumed that the slit had moved and a new time series was started at point $i + 1$. The sensitivity threshold D and the minimum number of cases N_0 are provided by the user. We determined empirically their value from a posteriori considerations on attained signal-to-noise ratios and spatial resolutions, and the trade-off among both parameters. In the cases under analysis in this work the achieved noise levels laid between 5×10^{-5} and 2×10^{-4} of the continuum level, for a spatial resolution better than 1 arc sec and approaching at times 0.6 arc secs².

5. Results

In Fig. 4 we show for illustration some of the observed Stokes I and V profiles (dotted lines) and the correspondent fits from the inversion (solid lines) overplotted. The panel covers the diversity of magnetic field strengths found in the quiet sun as seen through the Mn I lines: from the weak intensity regime (hG) to the strong regime (kG).

5.1. The complete set of profiles

Only 14% of the total amount of data had a S/N ratio high enough to be inverted reliably: around 500 profiles. In Fig. 5, we show the resulting magnetic field strength and longitudinal flux density distributions (defined as filling factor times the longitudinal field strength) from those inversions. For better description, we have fitted the histograms to an empirical distribution function of the form $N(x) = a \exp(-x/b) + c$, where a , b and c are positives constants and $N(x)$ represents the number of cases. The introduction of a uniform distribution (the constant c) is required

to fit the flat part of the histograms, corresponding to intermediate and high strength and flux density values. Otherwise, the left part of the histogram, associated to weak strength and flux density values respectively, is well fitted with an exponential function. In the case of the field strength distribution, the fitting values are $a=672$, $b=51.3$ G and $c=8.6$, while in the case of the flux density $a=465$, $b=6.3$ Mx/cm² and $c=1.7$. The full widths at half maximum are respectively $\text{FWHM}_B = 54$ G and $\text{FWHM}_{\text{Flux}} = 4.6$ Mx/cm², and the correspondent mean values are $\langle B \rangle = 635$ G and $\langle \text{Flux} \rangle = 13$ Mx/cm².

Although one expects mostly maxwellian-like probability distribution functions (PDF) (Dominguez Cerdeña et al. 2006b) for the field strength, because of the coarseness of the histogram bins we prefer to fit the data with an exponential for the weak fields plus a constant distribution for the strong fields. In spite of that empirical fit, the histogram shown in Fig. 5 should not be translated into that the most probable strength of the inverted profiles is zero G, but rather some value between $0 < B < 75$ G, included in the lower bin of the histogram. The main information the histograms show is that the weak field strength regime (hG) is more abundant or dominates the surface coverage in the quiet sun conditions (70%). Nevertheless, a considerable amount of cases, 30% of the inverted profiles, present magnetic strengths superior to 1000 G, i.e. in the strong field strength regime, and cannot be considered as absent.

5.2. Network versus Intranetwork

In order to study how the presence of network and intranetwork regions influence the results shown in the histograms of Fig. 5, we proceed to differentiate the profiles between these two regions. To this purpose, we have used the Fe I line (5538.52 nm), sitting next to the Mn line, to set a criterion for the classification of profiles into one or the other regions. We arbitrarily set the following condition for points to belong to the photospheric network: If the level of the normalized circular polarisation V/I in the Fe I line exceeds a threshold value (TV) of the continuum level we consider the point as part of the network. All other points are considered intranetwork. We have initially used two threshold values: $TV = 1 \times 10^{-3}$ and 2×10^{-3} . In Fig. 6, we show both distributions (for field strength and longitudinal flux density) for each one of the regions. The solid line represent a $TV = 1 \times 10^{-3}$ and the gray dotted line is for $TV = 2 \times 10^{-3}$. When using the upper threshold, $TV = 2 \times 10^{-3}$ corresponding to higher flux concentrations, the contribution of profiles in the weak strength regime in the network region has almost disappeared while the number of profiles with strong field strengths has been just slightly modified. On the other hand, the use of the moderate threshold value 1×10^{-3} already identifies the profiles in the strong regime in the network region as well as a non negligible contribution of fields with weak strengths. In the intranetwork set, with either one or the other threshold the general shape of the histogram remains unchanged. It is clear that a different threshold value able to radically change the general shapes of the histograms for both (network and intranetwork) regions would be unacceptably high and unrealistic. Therefore, and keeping in mind that small dependence of the network-assigned weak fields on the threshold value, we adopt in what follows the network and intranetwork definitions using the lower TV , with correspondent distributions function represented by the solid line in Fig. 6. To complete the justification on the criterion use for classification into network and intranetwork, let us add that from previous work (López Ariste et al. 2006) we know that the described classification, however arbitrary, successfully identifies most of

² The spatial resolution was gauged by the crude and simple method of counting and measuring the size of brighter-than-average and darker-than-average regions as visible in the continuum cuts along the slit.




Fig. 4. Examples of inverted profiles. The dotted lines correspond to the observed data and the solid line to the respective fitted profile. The profiles were inverted without any smoothing, but in this plot a 2-pixel smoothing has been applied to observed data.




Fig. 5. *Top panels:* Histograms of the distribution of the magnetic field strength and the longitudinal magnetic flux represented in logarithmic scale. *Bottom panels:* The same histograms in normal scale. The dashed lines represent the fitted distribution functions $N(x)$. The bin sizes are 75 G and 5 Mx/cm² in field strength and flux, respectively.

the network as identified visually over a magnetograph image. In terms of magnetic flux, the threshold value of 1×10^{-3} , would correspond roughly to 10 Mx/cm², and it is certainly biased towards including any flux concentration found in the intranetwork into the network, and conversely, any low flux region in the network will be included in the intranetwork dataset, a bias that we should remember in the conclusions.

After classification, from a sample of 477 inverted cases, it was found that 70% belonged to the intranetwork region, while the resting 30% were from the network. As already noticed in the previous paragraph, the correspondent distributions for the magnetic field strength in the network region, represented by the solid line in the left panel in Fig. 6, clearly differs from the inter-network distribution and also from the complete dataset distribution: It presents an apparent growth in the number of cases in the strong field regime, in a striking bimodal distribution. On the other side, the intranetwork distribution keeps a similar shape to the one found before: The intranetwork appears to be predominantly permeated by weak fields.

The main conclusion arising from these separate distributions is that the derived magnetic strength distribution from the inversions of the Mn I line in the visible range at 553 nm, in the low-flux quiet sun or intranetwork region (taking into account the known bias of our definition of intranetwork) is dominated in number of cases by weak field-

strength fields. This is in accord with previous results obtained with the analysis by measurement in the infrared range (Khomenko et al. 2003; Lin 1995; Lin & Rimmele 1999), in the visible (Keller et al. 1994) and with the recently retrieved distributions from the very high spectral resolution (0.3 arc sec) HINODE data (Orozco Suárez et al. 2007). Moreover, we found that the most probable expected strength is $B < 100$ G, close to the result obtained from simultaneous inversions from the infrared and from the visible recently published in Martínez González et al. (2008).

On the other hand, strong flux regions, mostly the photospheric network, rather show a bimodal distribution of fields with a contribution of weak field-strength fields comparable to that found in weak-flux regions, and an increased appearance of kG fields (Dominguez Cerdeña et al. 2006b; Lites 2002).

In the table 4 we summarise these results in terms of inversion percentages for different magnetic strengths as seen in the histograms of the analysed data sets previously discussed.

6. Conclusions

In this work we contribute to explore the diagnostic capabilities of Mn lines with strong hyperfine coupling for the magnetic topology of the quiet sun. We built an inversion code for the Mn line at 553.7 nm formed in a Milne-Eddington atmosphere, and




Fig. 6. Histograms of network and intranetwork regions. The solid (dotted) line correspond to a threshold value of 1×10^{-3} (2×10^{-3}). In the network, the contribution of profiles with strength superior to 1 kG is considerably more important than in the intranetwork.

Table 4. Comparison between the total set of inverted profiles and the Network-Intranetwork profiles.

	Total inverted	Intranetwork	Network
Number of Profiles	477	339	138
Strong regime (kG)	30%	17%	63%
Weak regime (hG)	70%	83%	37%
$\langle B \rangle$ (G)	635	435	1130
$\langle \text{Flux} \rangle$ (Mx/cm ²)	12.8	7.3	26.5

tested it under the conditions typical of the quiet sun in terms of signal amplitudes and noise levels. Such conditions are highly constraining: in most of the cases the magnetic field information is contained exclusively in the Stokes V profile, the linear polarization signals being under noise levels and the intensity profile being dominated by the non-magnetic portion of the resolution element. Under such restrictive conditions the number of independent observables for the magnetic field is highly reduced (Asensio Ramos et al. 2007) and it sets a limit to the number of free parameters of our model. Previous works have shown that the Mn lines increased the number of observables thanks to the spectral features appearing under certain magnetic regimes. The inversion code we tested in this work takes advantage of it and allows us to improve over the simplistic weak-field approximation of previous works into a more sophisticated Milne-Eddington atmosphere and yet to conserve the possibility of disentangling strong from weak fields.

Since most of the other parameters in the Milne-Eddington model, aside from field strength and flux, were not determined accurately under quiet-sun conditions, we doubt that any more sophisticated model can be used in the diagnostics at this point without the addition of further independent observables (e.g. with other spectral lines). However, more sophisticated models can shed light into implicit biases in such unrealistic picture of the solar photosphere as a Milne-Eddington atmosphere is, e.g. (Sánchez Almeida et al. 2008).

We have paid special attention throughout this work to the numerical tests made to ascertain whether, under present observational conditions of the quiet sun, the weak fields were disentangled from the strong fields for similar net fluxes. The answer is affirmative and therefore we proceeded to apply the code to real data taken with the THEMIS telescope under good seeing

conditions. Even in a telescope like THEMIS, focused on performing the most sensitive polarimetry possible, the acquisition of the data at the required signal-to-noise levels implied a trade-off in spatial resolution (seldomly better than 1 arc sec) that limits our results. We therefore constrained ourselves to separate the photospheric network from the intranetwork in our study, and restrained of identifying smaller scales, like granules and intergranules.

To identify network from intranetwork we applied a rather poor rule fixing at 10^{-3} the smallest amplitude of the Stokes V profile respect to the continuum intensity for a point to belong to the photospheric network. Such a rule obviously classifies as network any high flux concentration, mostly found over the network, but not totally absent from the intranetwork. Conversely it calls intranetwork any low flux concentrations, mostly found all over the intranetwork but not absent from the network either. With such bias in mind we decided to keep the names of network and intranetwork for the two classes because a small change in such threshold value did not change the general shape of the histograms, but also because of the following reasons.

After inversion, the intranetwork points show a distribution of field strengths whose dominant feature can be easily fitted with an exponential tail. We use the noun *tail* because an exponential distribution would result in a non-zero probability for null fields, something unacceptable for a vector field. Such a distribution, showing that the quiet sun regions are mostly permeated of weak strength fields, would be in full agreement with previous results in both Hanle and Zeeman effects (Manso Sainz et al. 2004; Lin & Rimmele 1999; Khomenko et al. 2003; Martínez González et al. 2008) and reveals a random or disorganised vector magnetic field at scales much smaller than present spatial resolution, if Hanle and Zeeman diagnostics are to be coherent with each other and with a field strength whose highest probability remains below 100G (Sánchez Almeida et al 2003).

The network points, on the other hand, show a bimodal distribution: the exponential tail found in the intranetwork distribution is still present, peaking at field strengths below 100G. But on top of that disorganised magnetic regime we observe the appearance of stronger fields, in the kiloGauss regime. This second distribution of fields would be in agreement with the presence of one or several concentration mechanisms, able to stabilise, organise and perhaps amplify the statistical fluctuations

of the ubiquitous turbulent field in those places where photospheric dynamics or sheer accidents are able to maintain a temporal coherence bigger than the typical magnetic diffusivities (Parker 1982). One can heuristically think that the strong downrafting plumes in the vertex of the convection cells are such places (Nordlund et al. 1992; Rast 2003), thus defining the magnetic photospheric network as a non-continuous succession of bright network points (Muller & Roudier 2003) where magnetic fields are organised in vertical structures and made stable by the downdrafting plasma. Other smaller plumes found in mesogranular or granular scales (Rast 2003) may also survive long enough to allow the statistical concentration of high fluxes, to be found here and there throughout the intranetwork and that, in our analysis, would have been included in the network group.

The cartoon just drawn above is in agreement with the histograms measured in the present work over the quiet sun. In particular, we are prone to the presence of an ubiquitous random field, as inferred from Hanle diagnostics, and whose statistical fluctuations can be seen as Zeeman signatures over deep magnetograms, but otherwise highly disorganized at present resolutions except when plasma movements create conditions, as around convection plumes, stable enough to concentrate and build up net magnetic flux. In any case, with the inversions performed here on the Mn I in the visible region of the spectra, at 553 nm, we have found that the distributions of the strength fields in the quiet Sun regions are dominated in occurrence by weak strength fields except in those areas related with the network.

Acknowledgements. JCRV thanks to M.J. Martínez González for all helpful discussions and also to the referee, J. Sánchez Almeida, for all the comments that served to improve the final state of the manuscript.

References

- Abt A., 1952, ApJ, 115, 199
 Asensio Ramos, A., Socas-Navarro, H., López Ariste, A., & Martínez González, M. J. 2007, ApJ, 660, 1690
 Bellot Rubio L. R. & Collados M., 2003, A&A, 406, 357
 Domínguez Cerdeña I., Sánchez Almeida J. & Kneer F., 2006, ApJ, 646, 1421
 Domínguez Cerdeña I., Sánchez Almeida J. & Kneer F., 2006b, ApJ, 636, 496
 Domínguez Cerdeña I., & Kneer F. & Sánchez Almeida J., 2003, ApJ, 582, L55
 Gelly B & López Ariste, A., 2006, sf2a.conf, 539
 Golub, G. H. & Van Loan, C. F., 1996, *Matrix Computations*, 3rd ed., Johns Hopkins University Press, Baltimore.
 Hansen, P. C., 1992, SIAM review, 34(4), 561
 Keller, C. U., Deubner, F.-L., Egger, U., Fleck, B., & Povel, H. P. 1994, A&A, 286, 626
 Khomekno E.V., Collados M., Solanki S., Lagg A. & Trujillo Bueno J., 2003, ApJ, 115, 199
 Lin, H. 1995, ApJ, 446, 421
 Lin, H. & Rimemele 1999, ApJ, 514, 448
 Lites B.W., 2000, ApJ, 573, 431
 López Ariste, A., Rayrole, J., & Semel, M. 2000, A&A, 142, 137
 López Ariste A. & Casini R. 2002, ApJ, 580, 519
 López Ariste A. & Casini R. 2002, ApJ, 575, 529
 López Ariste A., Tomczyk S., & Casini R., 2006, A&A, 454, 663
 López Ariste A, Martínez González M.J. & Ramírez Vélez J.C., 2006, A&A, 464, 351
 López Ariste, A., & Semel, M. 1999, A&A, 139, 417
 Manso Sainz R., Land Degl'Innocenti L. & Trujillo Bueno J. ApJ, 2004, 614, 89
 Martínez González, M. J., Collados, M., & Ruiz Cobo, B. 2006, A&A, 456, 1159
 Martínez Gonzalez M.J., Collados M., Ruiz Cobos B. & Beck C., 2008, A&A, 477, 953
 Muller R & Roudier Th., SoPh, 1992, 141 27
 Nordlund A., Brandenburg A., Jennings R.L., Rieutord M., Ruokolainen J., Stein R.F. & Tuominen I. 1992, ApJ, 392, 647
 Orozco Suárez, D., et al. 2007, ApJ, 670, L61
 Parker E.N., 1982, ApJ, 256, 292
 Kurucz R.L. 1993, Phy. Scr., T47, 110
 Rast, M.P., 2003, ApJ, 597, 1200
 Rees, D. E., López Ariste, A., Thatcher, J., & Semel, M. 2000, A&A, 355, 759
 Sánchez Almeida, J., & Landi degl'Innocenti, E., 1996, Sol. Phys., 164, 203
 Sánchez Almeida, J., & Lites, B. W. 2000, ApJ, 532, 1215
 Sánchez Almeida, J., Emonet T., & Cattaneo F. 2003, ApJ, 585, 536
 Sánchez Almeida, J., Vitićchié, B., Landi Degl'Innocenti, E., & Berrilli, F., 2008, ApJ, 675, 906
 Socas Navarro H., López Ariste A. & Lites B.W., 2001, ApJ, 553, 949
 Socas Navarro, H., & Sánchez Almeida, J. 2003, ApJ, 593, 581
 Stenflo J.O., 1982, SoPh, 80, 209S